Tables in OAR 437-002-0156

Table reference in (5)(e)(A) in High Heat Practices section

Table 1. Minimum employer-specific rest break durations and intervals:

Heat index temperature (° F)	Rest break durations	
90 or greater	10 minutes every two hours	
100 or greater	15 minutes every hour	

- (i) The effect of personal protective equipment (PPE) on the body's ability to retain heat;
- (ii) The effect of the type of work clothing on the body's ability to retain heat;
- (iii) Relative humidity, whether work activities are indoors or outdoors; and
- (iv) The intensity of the work being performed.

Table reference in (5)(e)(C) in High Heat Practices section

Table 2. Minimum simplified rest break durations and intervals:

Heat index temperature (° F)	Rest break durations
90 or greater	10 minutes every two hours
95 or greater	20 minutes every hour
100 or greater	30 minutes every hour
105 or greater	40 minutes every hour

Note: The Table 2 work/rest schedule is only required during the specified heat index temperatures.

OAR 437-002-0156 Appendix A

Information for Heat Illness Prevention

To protect the health and safety of employees from heat-related illnesses, employers should consider using the resources below. Please note that some resources may use temperatures in Celsius instead of Fahrenheit. To convert to degrees Fahrenheit, use this formula: Fahrenheit ($^{\circ}$ F) = (Celsius x 1.8) + 32

1. Most heat-related illnesses affect workers who do strenuous physical activity. When workers engage in intense work, their bodies create heat. This "metabolic" heat combines with environmental heat (from temperature, sunlight, humidity, etc.) so workers' core temperature can rise to dangerous levels. To prevent a hazardous combination of environmental and metabolic heat, employers should be aware of workers' activity level. Workload can be classified as rest, light, moderate, heavy, or very heavy. See information provided in the tab on Metabolic Heat and Workload (Physical Activity level) for additional detail on estimating worker's workload.

Table 1.Metabolic Heat and Workload (Physical Activity Level)

Level of Workload / Physical Activity *	Examples	Metabolic Rate in Watts, "typical" recognizing that different ways of doing
		the same task may lead to dramatically different
		wattage
Rest	Sitting	115
	 Thinking 	
Light	 Sitting with minimal hand and arm work 	180
	Sewing	
	 Writing or drawing 	
	Driving a car	
	 Occasional or slow walking 	
	 Stooping, crouching, or kneeling 	
	 Standing watch 	

^{*} Workers who are overweight or obese might produce more metabolic heat than other workers who perform the same tasks. The above table assumes a 70-kg (154-pound) worker.

Table 1 was partially recreated from federal OSHA's guidance on Heat Hazard recognition, which can be accessed at: https://www.osha.gov/heat-exposure/hazards.

2. The OSHA-NIOSH Heat Safety Tool app is a useful resource for planning outdoor work activities based on how hot it feels throughout the day. It features real-time heat index and hourly forecasts specific to your location, as well as occupational safety and health recommendations from OSHA and NIOSH. It can be accessed and downloaded at: https://www.osha.gov/heat/heat-app

3. Work/rest schedules.

A. Table 2. Work/rest schedules for workers wearing normal work clothing*

Adjusted	Light work	Moderate work	Heavy work
temperature (°F) [†]	(minutes work/rest)	(minutes work/rest)	(minutes work/rest)
90	Normal	Normal	Normal
91	Normal	Normal	Normal
92	Normal	Normal	Normal
93	Normal	Normal	Normal
94	Normal	Normal	Normal
95	Normal	Normal	45/15
96	Normal	Normal	45/15
97	Normal	Normal	40/20
98	Normal	Normal	35/25
99	Normal	Normal	35/25
100	Normal	45/15	30/30
101	Normal	40/20	30/30
102	Normal	35/25	25/35
103	Normal	30/30	20/40
104	Normal	30/30	20/40
105	Normal	25/35	15/45
106	45/15	20/40	Caution [‡]
107	40/20	15/45	Caution [‡]
108	35/25	Caution [‡]	Caution [‡]
109	30/30	Caution [‡]	Caution [‡]
110	15/45	Caution [‡]	Caution [‡]
111	Caution [‡]	Caution [‡]	Caution [‡]
112	Caution [‡]	Caution [‡]	Caution [‡]

^{*}With the assumption that workers are physically fit, well-rested, fully hydrated, under age 40, and have adequate water intake and that there is 30% RH [relative humidity] and natural ventilation with perceptible air movement.

Full sun (no clouds): Add 13° Partly cloudy/overcast: Add 7°

No shadows visible/work is in the shade or at night: no adjustment

Per relative humidity:

10%: Subtract 8° 20%: Subtract 4° 30%: No adjustment

40%: Add 3° 50%: Add 6° 60%: Add 9°

Adapted from EPA [1993]

[†]Note: Adjust the temperature reading as follows before going to the temperature column in the table:

[‡]High levels of heat stress; consider rescheduling activities.

Table 2 above was recreated from the following publication; see page 76 in NIOSH [2016]. NIOSH criteria for a recommended standard: occupational exposure to heat and hot environments. By Jacklitsch B, Williams WJ, Musolin K, Coca A, Kim J-H, Turner N. Cincinnati, OH: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health, DHHS (NIOSH) Publication 2016-106.

B. Table 3. Work/rest schedules for those wearing chemical resistant suits.

		Light work		Moderate work			Heavy work		
Air Temp (°F)	Full sun	Partly cloudy	No sun [†]	Full sun	Partly cloudy	No sun [†]	Full sun	Partly cloudy	No sun [†]
75	Normal	Normal	Normal	Normal	Normal	Normal	35/25 [‡]	Normal	Normal
80	30/30	Normal	Normal	20/40	Normal	Normal	10/50	40/20	Normal
85	15/45	40/20	Normal	10/50	25/35	Normal	Caution §	15/45	40/20
90	Caution §	15/45	40/20	Caution §	Caution §	25/35	Stop work	Caution§	15/45
95	Stop	Stop	15/45	Stop	Stop	Stop	Stop	Stop	Stop
	work	work		work	work	work	work	work	work

^{*}With the assumption that workers are heat-acclimatized, under the age of 40, physically fit, well-rested, fully hydrated, and wearing Tyvek coveralls, gloves, boots, and a respirator. Cooling vests may enable workers to work for longer periods. Adjustments must be made when additional protective gear is worn.

Adapted from EPA [1993]

Table 3 above was recreated from the following publication; see page 77 in NIOSH [2016]. *NIOSH criteria for a recommended standard: occupational exposure to heat and hot environments.* By Jacklitsch B, Williams WJ, Musolin K, Coca A, Kim J-H, Turner N. Cincinnati, OH: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health, DHHS (NIOSH) Publication 2016-106.

4. Acclimatization.

Table 4. Acclimatization in workers

Topics	Additional information			
Disadvantages of	 Readily show signs of heat stress when 			
being unacclimatized	exposed to hot environments.			
	 Difficulty replacing all of the water lost in 			
	sweat.			
	Failure to replace the water lost will slow			
	or prevent acclimatization.			
Benefits of acclimatization	 Increased sweating efficiency (earlier 			
	onset of sweating, greater sweat			

[†]No shadows are visible or work is in the shade or at night.

[‡]35 minutes work and 25 minutes rest each hour.

[§]High levels of heat stress; consider rescheduling activities.

Acclimatization plan	production, and reduced electrolyte loss in sweat). Stabilization of the circulation. Work is performed with lower core temperature and heart rate. Increased skin blood flow at a given core temperature. Gradually increase exposure time in hot environmental conditions over a period of 7 to 14 days. For new workers, the schedule should be no more than 20% of the usual duration of work in the hot environment on day 1 and a no more than 20% increase on each additional day. For workers who have had previous experience with the job, the acclimatization regimen should be no more than 50% of the usual duration of work in the hot environment on day 1, 60% on day 2, 80% on day 3, and 100% on day 4. The time required for non—physically fit individuals to develop acclimatization is about 50% greater than for the physically fit.
Level of acclimatization	Relative to the initial level of physical fitness and the total heat stress experienced by the individual.
Maintaining acclimatization	 Can be maintained for a few days of nonheat exposure. Absence from work in the heat for a week or more results in a significant loss in the beneficial adaptations leading to an increased likelihood of acute dehydration, illness, or fatigue. Can be regained in 2 to 3 days upon return to a hot job. Appears to be better maintained by those who are physically fit. Seasonal shifts in temperatures may result in difficulties. Working in hot, humid environments provides adaptive benefits that also apply

in hot, desert environments, and vice				
versa.				
 Air conditioning will not affect 				
acclimatization.				

Adapted from [Moseley 1994; Armstrong and Stoppani 2002; DOD 2003; Casa et al. 2009; ACGIH 2014; OSHA-NIOSH 2011].

Table 4 above was recreated from the following publication; see page 34. NIOSH [2016]. NIOSH criteria for a recommended standard: occupational exposure to heat and hot environments. By Jacklitsch B, Williams WJ, Musolin K, Coca A, Kim J-H, Turner N. Cincinnati, OH: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health, DHHS (NIOSH) Publication 2016-106.

5. Clothing adjustment factors.

Table 5. Clothing adjustment factors exist for various types of clothing.

Clothing adjustment factors (°C-WBGT)

Clothing	Previous	2006
Work clothing (baseline)	0	0
Cloth coveralls	3.5	0
Double-layer cloth clothing	5	3
Spunbound melt-blown synthetic (SMS) coveralls	-	0.5
Polyolefin coveralls	-	1
Limited-use vapor-barrier coveralls	_	11

Adapted from Bernard TE, Threshold Limit Values for Physical Agents Committee, ACGIH [2014].

Table 5 above was recreated from the following publication; see page 19. NIOSH [2016]. NIOSH criteria for a recommended standard: occupational exposure to heat and hot environments. By Jacklitsch B, Williams WJ, Musolin K, Coca A, Kim J-H, Turner N. Cincinnati, OH: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health, DHHS (NIOSH) Publication 2016-106.